



## CHAIRSIDE IMPLANT SERVICES

*We Come To You...Chairside!*

631-581-5121

### **PRE-OP INSTRUCTIONS FOR IMPLANT PLACEMENT**

- Premedicate with four, 500mg tablets (2000mg) of Amoxicillin 1 hour before your implant placement, then take 1 tablet (500mg) every 8 hours for 7 days.
- If you are allergic to Amoxicillin, or Penicillin, premedicate with Clindamycin instead. Take four, 150 mg tablets (600mg) 1 hour before your implant placement and then take 1 tablet (150mg) every 8 hours for 7 days.
- If you take aspirin on a daily basis, stop taking it for 14 days prior to your implant placement. Consult with your physician prior to stopping the aspirin.
- 400mg of ibuprofen (Advil or Motrin) is recommended 1 hour prior to implant placement
- Wear comfortable clothing. Single implant procedures can take up to 1 hour. Multiple implant procedures can take 2 hours.
- For full-arch implant procedures, you will be on a soft diet for 7-14 days after implant placement. Juice, soup, yogurt and jello will be advised for the first 24-48 hours. Then soft foods can be added gradually.
- For single tooth or sectional arch implant procedures, you will be instructed to chew on the opposite side for a period of time.
- Bring your dentures, stents and/or temporary bridges to the office the day of implant placement.
- Most patients experience no pain during or after the procedure. Minor discomfort is to be expected for multiple implant placement.